

Female Breast Cancer Deaths, by Age and Race, Maine, 2004-2008

	Count	Crude Rate		Age-adjusted Rate	
		Rate	95% CI	Rate	95% CI
Age					
00-14 years	NSD	NSD	NSD	NApp	NApp
15-19 years	NSD	NSD	NSD	NApp	NApp
20-24 years	59	0.1	0.1 - 0.1	NApp	NApp
25-29 years	355	0.7	0.6 - 0.8	NApp	NApp
30-34 years	1,432	2.9	2.8 - 3.1	NApp	NApp
35-39 years	3,877	7.5	7.2 - 7.7	NApp	NApp
40-44 years	7,773	13.9	13.6 - 14.2	NApp	NApp
45-49 years	12,822	22.5	22.1 - 22.9	NApp	NApp
50-54 years	17,520	33.6	33.1 - 34.1	NApp	NApp
55-59 years	21,449	47.0	46.3 - 47.6	NApp	NApp
60-64 years	21,704	60.7	59.9 - 61.5	NApp	NApp
65-69 years	20,281	72.4	71.4 - 73.4	NApp	NApp
70-74 years	20,202	85.1	83.9 - 86.3	NApp	NApp
80-84 years	22,922	130.6	129.0 - 132.3	NApp	NApp
85+ years	31,150	177.6	175.7 - 179.6	NApp	NApp
Race					
White	1,966	23.2	22.2 - 24.3	30.3	28.9 - 31.6
Black	NSD	NSD	NSD	NSD	NSD
Other*	16	22.8	12.0 - 38.9	14.6	8.3 - 23.7

Data Source: Surveillance, Epidemiology, and End Results SEER Program, National Cancer Institute, Underlying mortality data provided by National Center for Health Statistics. Rates are calculated using SEER*Stat Version 7.0.5.

Cancer Deaths: Deaths with malignant cancer as the underlying cause of death.

Female Breast Cancer: SEER Cause of Death Recode: 26000 (which include 1999-2009: ICD-10 codes C50; 1993-1998: ICD-9 codes 174-175).

Crude rates are deaths per 100,000 population.

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.

95% CI: 95% confidence interval of the rate.

*Other Race includes American Indian/AK Native and Asian/Pacific Islander

NSD = Non sufficient data. Data not available due to too few cases.

NApp: Not applicable age-adjusted rates are not calculated for age-specific groups.

